



UNIVERSITÀ
DI PAVIA




Istituto Nazionale di Fisica Nucleare


On the way to transversity: A PhD triathlon

PhD candidate:
Luca Polano

Supervisor:
Marco Radici
Alessandro Bacchetta

End of year seminars 2025






Quarks &
gluons




Depends on the energy scale



High energies






Quarks &
gluons

High energies


Low, $\Lambda_{QCD} \sim 200 \text{ MeV}$




$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$




$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$



$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$




$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$



$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$


PDF



$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$

PDF


f_1



$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$

PDF


g_1



$$At \sim \Lambda_{QCD} \sim 200 \text{ MeV}$$


PDF

h_1



Why caring about transversity?

- Have a better knowledge of the proton structure



Why caring about transversity?

- Have a better knowledge of the proton structure
- Can be used to investigate physics beyond the Standard Model

Chiral-odd structures do not appear in the SM
Tree level Lagrangian


Neutrino β -decay

$$g_T = \delta u - \delta d$$

$$\delta q = \int_0^1 dx h_1^q(x)$$

constraints on CP violation

N.Sato et al, Phys. Rev. D 109, (2024) 034024



Why caring about transversity?

- Have a better knowledge of the proton structure
- Can be used to investigate physics beyond the Standard Model


Chiral-odd structures do not appear in the SM
Tree level Lagrangian



Neutrino β -decay


$$g_T = \delta u - \delta d$$

$$\delta q = \int_0^1 dx h_1^q(x)$$

constraints on CP violation











X












Fragmentation Functions




Fragmentation Functions






Fragmentation Functions









Fragmentation Functions












$$x_B \leftrightarrow z$$


D_1

Electron-positron annihilation




Electron-positron annihilation

H_1^Δ





Semi-inclusive Deep Inelastic Scattering

h_1




Semi-inclusive Deep Inelastic Scattering





$$\frac{d\sigma}{dx dQ^2 \dots} = H \otimes PDF \otimes FF$$


Semi-inclusive Deep Inelastic Scattering



Ex.: NLO


$$\frac{d\sigma}{dx dQ^2 \dots} = H \otimes PDF \otimes FF$$

2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{\leq, q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$


2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{\leq, q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$

How to access
them?


2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{\leq, q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$




2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{4,q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$




2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{\leq,q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$




2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{\leq,q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$




2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{4,q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$



2 measured hadrons + X



$$A_{UT}^{hh} \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B) \cdot H_1^{\leq,q}(z, M_h)}{\sum_q e_q^2 \cdot f_1^q(x_B) \cdot D_1^q(z, M_h)}$$



GOALS of the D_1 extraction

GOALS of the D_1 extraction

- 2017 BELLE data of $e^+e^- \rightarrow \pi^+\pi^-X$ at $\sqrt{S} = 10.58 \text{ GeV}$

GOALS of the D_1 extraction

- 2017 BELLE data of $e^+e^- \rightarrow \pi^+\pi^-X$ at $\sqrt{S} = 10.58 \text{ GeV}$
- Use of Monte-Carlo simulation for flavor separation only


GOALS of the D_1 extraction

- 2017 BELLE data of $e^+e^- \rightarrow \pi^+\pi^-X$ at $\sqrt{S} = 10.58 \text{ GeV}$
- Use of Monte-Carlo simulation for flavor separation only
- Extend the analysis up to NNLO

GOALS of the D_1 extraction

- 2017 BELLE data of $e^+e^- \rightarrow \pi^+\pi^-X$ at $\sqrt{S} = 10.58 \text{ GeV}$
- Use of Monte-Carlo simulation for flavor separation only
- Extend the analysis up to NNLO
- Explore a Neural Network parameterisation


PHYSICS INFORMED



71 par




PHYSICS INFORMED



71 par


NEURAL NETWORK




NN architecture

$[2, 25, 5] \rightsquigarrow$ 205 par


PHYSICS
INFORMED



71 par




NEURAL
NETWORK




NN architecture

[2,25,5] $\rightsquigarrow$ 205 par


PHYSICS
INFORMED



71 par




NEURAL
NETWORK




NN architecture

[2,25,5] $\rightsquigarrow$ 205 par


PHYSICS INFORMED



71 par






NEURAL NETWORK



NN architecture

[2,25,5] $\rightsquigarrow$ 205 par

Predictions: Physics informed



NNLO

100 replicas

$$\chi_u^2 = 0.380$$





$$\chi_d^2 = 0.613$$

$$\chi_s^2 = 0.876$$

$$\chi_c^2 = 0.879$$

$$\chi_{tot}^2 = 0.687$$

Predictions: Neural Network



NNLO

100 replicas

$$\chi_u^2 = 0.209$$


$$\chi_d^2 = 0.530$$

$$\chi_s^2 = 0.793$$




$$\chi_c^2 = 0.606$$

$$\chi_{tot}^2 = 0.535$$




Comparison



NN seems to describe data better




NN has less sensitivity on resonances



DI-HADRON FF


Physics informed



$u=d$


ratio offset $\Delta = 0.5$

Δ



DI-HADRON FF


Physics informed




$u=d$

ratio offset $\Delta = 0.5$

Δ




DI-HADRON FF Neural Network




$u \neq d$
ratio offset $\Delta = 2.5$


Δ




DI-HADRON FF Neural Network



$u \neq d$
ratio offset $\Delta = 2.5$



Comparison at NNLO:









Similar features and compatible results


NN has wider uncertainty bands

Resonances are better captured by the P.I. fit



Gluon?




Gluon and up bands at NNLO



$z = 0.50$



$z = 0.50$



CONCLUSIONS AND OUTLOOKS

CONCLUSIONS AND OUTLOOKS

- **Good agreement for both fits**

CONCLUSIONS AND OUTLOOKS

- Good agreement for both fits
- Hierarchy of quark flavours (charm?)

CONCLUSIONS AND OUTLOOKS

- Good agreement for both fits
- Hierarchy of quark flavours (charm?)
- Gluon unconstrained

CONCLUSIONS AND OUTLOOKS

- Good agreement for both fits
- Hierarchy of quark flavours (charm?)
- Gluon unconstrained
- Use D_1 to extract $H_1^{\lessdot}$ and then h_1

CONCLUSIONS AND OUTLOOKS



A cyclist in a red jersey is riding a road bike on a paved road. The background shows a clear sky and distant hills. A red dotted arrow points from the text 'CONCLUSIONS AND OUTLOOKS' to the road surface.

CONCLUSIONS AND OUTLOOKS

- Good agreement for both fits



CONCLUSIONS AND OUTLOOKS

- Good agreement for both fits
- Hierarchy of quark flavours (charm?)



CONCLUSIONS AND OUTLOOKS

- Good agreement for both fits
- Hierarchy of quark flavours (charm?)
- Gluon unconstrained




CONCLUSIONS AND OUTLOOKS

- Good agreement for both fits
- Hierarchy of quark flavours (charm?)
- Gluon unconstrained
- Use D_1 to extract $H_1^{\triangleleft}$ and then h_1




1 measured hadron+ X



$$A_{UT}^h \sim \frac{\sum_q e_q^2 \cdot h_1^q(x_B, k_T) \otimes H_1^{\perp, q}(z, p_{\perp})}{\sum_q e_q^2 f_1^q(x_B, k_T) \otimes D_1^q(z, p_{\perp})}$$

2012 extraction by Pavia group



A.Bacchetta, M.Radici, A.Bianconi, A.Courtoy, Phys. Rev. D 85, (2012) 114023

Fit of MonteCarlo simulation

79 free parameters

LO

2017 BELLE data of $e^+e^- \rightarrow \pi^+\pi^-X$ at $\sqrt{S} = 10.58$ GeV


Latest extraction:

+ MonteCarlo simulation

LO

u
 s
 c
 b
 g


N.Sato et al, Phys. Rev. D 109, (2024) 034024



Flavour analysis

Physical review D 96 (2017)
R.Siedl et al

$$\frac{d\sigma}{dzdM_hdQ^2} = \frac{4\pi\alpha^2}{Q^2} \sum_q e_q^2 D_1^q(z, Mh, Q) =$$



Flavour analysis

$$\frac{d\sigma}{dzdM_hdQ^2} = \frac{4\pi\alpha^2}{Q^2} \sum_q e_q^2 D_1^q(z, M_h, Q) = \frac{4\pi\alpha^2}{Q^2} \mathbf{D}(\mathbf{z}, \mathbf{M}_h, \mathbf{Q})$$

Need of extra information  Monte Carlo

$$\mathbf{R}_u^{\text{MC}} = \frac{\mathbf{D}_{\text{MC}}^u(\mathbf{z}, \mathbf{M}_h, \mathbf{Q})}{\mathbf{D}_{\text{MC}}(\mathbf{z}, \mathbf{M}_h, \mathbf{Q})}$$

 Build 4 pseudo-dataset


$$\mathcal{F}^u(z, M_h; Q^2) = \frac{4\pi\alpha^2}{Q^2} \mathbf{D}(\mathbf{z}, \mathbf{M}_h, \mathbf{Q}) \times \frac{\mathbf{D}_{\text{MC}}^u(\mathbf{z}, \mathbf{M}_h, \mathbf{Q})}{\mathbf{D}_{\text{MC}}(\mathbf{z}, \mathbf{M}_h, \mathbf{Q})}$$

$$\mathcal{F}^q$$


$$q = u, d, s, c$$

DI-HADRON FF

Physics informed



u=d

gluon  assumptions

$$D_1^g(z, M_h; Q^2) = N z^{\alpha_1} (1 - z)^{1+\alpha_2} \cdot D_1^u(z, M_h; Q^2)$$

N random unif. in (0,2)

α_1, α_2 random unif. in (0,1)